

Entangling gates in even Euclidean lattices such as the Leech lattice

Michel Planat¹

¹ Institut FEMTO-ST, CNRS, 32 Avenue de l'Observatoire, F-25044 Besançon, France.

Abstract. The group of automorphisms of Euclidean (embedded in $\mathbb{R}^n$) dense lattices such as the root lattices D_4 and E_8 , the Barnes-Wall lattice BW_{16} , the unimodular lattice D_{12}^+ and the Leech lattice Λ_{24} may be generated by entangled quantum gates of the corresponding dimension. These (real) gates/lattices are useful for quantum error correction: for instance, the two and four-qubit real Clifford groups are the automorphism groups of the lattices D_4 and BW_{16} , respectively, and the three-qubit real Clifford group is maximal in the Weyl group $W(E_8)$. Technically, the automorphism group $\text{Aut}(\Lambda)$ of the lattice Λ is the set of orthogonal matrices B such that, following the conjugation action by the generating matrix of the lattice, the output matrix is unimodular (of determinant ± 1 , with integer entries). When the degree n is equal to the number of basis elements of Λ , then $\text{Aut}(\Lambda)$ also acts on basis vectors and is generated with matrices B such that the sum of squared entries in a row is one, i.e. B may be seen as a quantum gate. For the dense lattices listed above, maximal multipartite entanglement arises. In particular, one finds a balanced tripartite entanglement in E_8 (the two- and three- tangles have equal magnitude $1/4$) and a GHZ type entanglement in BW_{16} . In this paper, we also investigate the entangled gates from D_{12}^+ and Λ_{24} , by seeing them as systems coupling a qutrit to two- and three-qubits, respectively. Apart from quantum computing, the work may be related to particle physics in the spirit of [1].

PACS numbers: 03.65.Ud, 03.67.Pp, 02.20.-a

1. Introduction

Entanglement is at the heart of many paradoxes of quantum mechanics and is also a useful resource in quantum information processing. In particular, entangled states play a key role in quantum error correction and in quantum computational speedup. The characterization of entanglement of a general multipartite state remains a mathematical challenge and there exist several inequivalent measures. A comprehensive account of the most important mathematical concepts used for quantifying and manipulating entangled states may be found in [2].

Some properties of entangled states may also be approached from differential geometry (two- and three-qubit Hilbert space geometry corresponds to Hopf fibrations, that are entanglement sensitive [3]), from topological spaces (for instance the 3-qubit

GHZ state corresponds to a borromean ring [4]), from finite geometries (by using the hyperplanes of the relevant generalized quadrangles [5] or in the black-hole analogy [6]. Besides, group theoretical concepts have been repetitively put forward for describing classical [7] and quantum error correction [8]. In this paper, we exploit the unnoticed but organic link between real entangling gates of quantum computation and the group of automorphisms of some highly symmetric even Euclidean lattices.

Let us define an n -dimensional Euclidean lattice Λ as a discrete additive subgroup of the real vector space $\mathbb{R}^n$, endowed with the standard Euclidean product, and spanned by a generator matrix M with rows in $\mathbb{R}^n$. The automorphism group $\text{Aut}(\Lambda)$ is the set of orthogonal matrices B such that under the conjugation action by the generating matrix M , one gets a unimodular matrix $U = MBM^{-1}$, that is (i) $\det U = \pm 1$ and (ii) U is an integer matrix (see [12], p. 90). When the degree n is equal to the number of basis elements of Λ , then $\text{Aut}(\Lambda)$ also acts on basis vectors and is generated with matrices B such that the sum of squared entries in a row is one, i.e. B may be seen as a quantum gate.

It will be shown that for a suitable choice of even Euclidean lattices Λ such as $\mathbb{Z}^n$ lattices, the root lattices D_4 and E_8 , the Barnes-Wall lattice Λ_{16} , the lattice D_{12}^+ and the Leech lattice Λ_{24} , one observes that the orthogonal group $\text{Aut}(\Lambda)$ is a group of entangling *quantum gates*. It can be considered as acting on a two-, three-, four-, a two-qubit/qutrit and a three-qubit/qutrit system, respectively. In retrospect, such lattices bear out our recent work relating D_4 and E_8 Weyl groups to entangled states arising from the joint basis of specific mutually commuting set of generalized Pauli observables [1, 11].

2. Entanglement arising from $\mathbb{Z}^n$ lattices

To illustrate our topic, let us start with the $\mathbb{Z}^n$ lattice whose lattice points are the integers and whose basis M is the identity matrix. The elementary cell is the n -dimensional hypercube and the automorphism group is a wreath product [14]

$$\text{Aut}(\mathbb{Z}^n) \cong \mathbb{Z}_2 \wr S_n = \mathbb{Z}_2^n \rtimes S_n, \quad (1)$$

of order $2^n n!$, where the symbol $\rtimes$ denotes the semidirect product and the wreath product $\wr$ corresponds to a permutation action of the symmetric group S_n on the n copies of the two-letter group $\mathbb{Z}_2$.

It is straightforward to check that the automorphism group $\text{Aut}(\mathbb{Z}^4)$ of the four-dimensional lattice contains the gate $\text{CNOT} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$, that is well known to

create two-qubit entanglement. More precisely, taking the input qubit states as $|0\rangle$ and $|\psi\rangle = a|0\rangle + b|1\rangle$, the action $\text{CNOT}(|0\rangle|\psi\rangle)$ ends up into the non-separable Bell state $a|00\rangle + b|11\rangle$. As it is well known, the CNOT gate is a primitive of universal quantum computing.

It has been found [1] that the entanglement arising from the two-qubit Pauli group is encapsulated in the following Mermin's square

$$\begin{array}{ccc} \sigma_x \otimes \sigma_x & \sigma_y \otimes \sigma_y & \sigma_z \otimes \sigma_z \\ \sigma_y \otimes \sigma_z & \sigma_x \otimes \sigma_z & \sigma_x \otimes \sigma_y \\ \sigma_z \otimes \sigma_y & \sigma_z \otimes \sigma_x & \sigma_y \otimes \sigma_x, \end{array} \quad (2)$$

where σ_x , σ_y and σ_z are the ordinary Pauli spin matrices. Every row and every column in the above square consists of mutually commuting operators sharing a basis of entangled states. The Mermin's square may be used for a proof of the Kochen-Specker theorem in a four-dimensional space by observing that the algebraic relations for the eigenvalues ± 1 of the observables contradicts the one for the eigenstates [1, 9].

Let us denote $r = \{\sigma_x \otimes \sigma_x, \sigma_y \otimes \sigma_y, \sigma_z \otimes \sigma_z\}$ and $s = \{\sigma_x \otimes \sigma_z, \sigma_z \otimes \sigma_x, \sigma_y \otimes \sigma_y\}$ the first row and the r.h.s. column in (2), the only ones that possess real entries. One may use the common eigenstate basis $M_r = \{(1, 0, 0, 1), (1, 0, 0, -1), (0, 1, 1, 0), (0, 1, -1, 0)\}$ of the set r for generating a lattice isometric to $\mathbb{Z}^4$. The natural action of the automorphism group on the basis M_r reads $\ddagger$.

$$\text{Aut}(\mathbb{Z}^4) = \left\langle S = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & 1 \\ 1 & 1 & 1 & -1 \\ 1 & -1 & -1 & -1 \\ 1 & 1 & -1 & 1 \end{pmatrix}, S' = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \end{pmatrix} \right\rangle, \quad (3)$$

It is noticeable that the rows of the generators S and S' of $\text{Aut}(\Lambda)$ encode the eigenstates of the triples of observables s and $\{\sigma_x \otimes 1_2, 1_2 \otimes \sigma_x, \sigma_x \otimes \sigma_x\}$, respectively (a partial list of the eigenstates, associated to the maximal sets of mutually commuting operators in the two-qubit Pauli group, may be found in [10]).

Then, following a similar reasoning, we construct the $\mathbb{Z}^8$ lattice by means of a generating basis built from the eigenstates of the following triple of three-qubit observables

$$s_3 = \sigma_z \otimes \{\sigma_x \otimes \sigma_z, \sigma_z \otimes \sigma_x, \sigma_y \otimes \sigma_y\}, \quad (4)$$

that follows from s by applying the left tensor product σ_z . The basis we select is the matrix S_3 first introduced in [1] (Eq. (9)). As for the two-qubit case above, the natural action of the automorphism group on the basis is obtained as

$$\text{Aut}(\mathbb{Z}^8) = \langle g_1, g_2 \rangle,$$

$\ddagger$ The reader may use the following commands in Magma [13] to check the calculations: “Mr:=Matrix(4,4,[1,0,0,1, 1,0,0,-1, 0,1,1,0, 0,1,-1,0]); L:=LatticeWithBasis(Mr); aut:=AutomorphismGroup(L:NaturalAction); L,aut;”

$$\text{with } g_1 = \frac{1}{2} \begin{pmatrix} . & . & 1 & 1 & . & 1 & . & 1 \\ . & . & 1 & 1 & . & -1 & . & -1 \\ -1 & 1 & . & . & 1 & . & -1 & . \\ 1 & -1 & . & . & 1 & . & -1 & . \\ 1 & 1 & . & . & . & 1 & . & -1 \\ 1 & 1 & . & . & . & -1 & . & 1 \\ . & . & -1 & 1 & 1 & . & 1 & . \\ . & . & 1 & -1 & 1 & . & 1 & . \end{pmatrix}. \quad (5)$$

The matrix g_2 is not made explicit but it has a similar form and properties as g_1 . Let us now investigate the type of entanglement contained in the 3-qubit generators/gates g_1 and g_2 . We single out the state encoded by the fourth row of g_1

$$|\psi\rangle = \frac{1}{2}(|000\rangle - |001\rangle + |100\rangle - |110\rangle). \quad (6)$$

A quantitative measure of tripartite entanglement is the 3-tangle [15, 16]

$$\tau^{(3)} = 4 |(T^{001} - T^{000})^2 - 4P_{B_1}^{00}P_{B_0}^{00}|, \quad (7)$$

which contains the four determinants

$$T^{000} = \det \begin{pmatrix} a_{000} & a_{011} \\ a_{100} & a_{111} \end{pmatrix}, T^{001} = \det \begin{pmatrix} a_{001} & a_{010} \\ a_{101} & a_{110} \end{pmatrix},$$

$$P_{B_0}^{00} = \det \begin{pmatrix} a_{000} & a_{001} \\ a_{100} & a_{101} \end{pmatrix} \text{ and } P_{B_1}^{00} = \det \begin{pmatrix} a_{010} & a_{011} \\ a_{110} & a_{111} \end{pmatrix}.$$

To quantify bipartite entanglement the strategy is as follows [16]. Take the density matrix $|\psi\rangle\langle\psi|$ of the 3-qubit system and trace out over the bipartite subsystems to obtain a reduced density matrix ρ . A measure of two-qubit entanglement is the tangle $\tau = C^2$, with the concurrence $C(\rho) = \max\{0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}\}$, where the λ_i are non-negative eigenvalues of the product $\rho\tilde{\rho}$ arranged in decreasing order, $\tilde{\rho} = (\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$ is the spin-flipped density matrix and $*$ denotes the complex conjugate.

Two important families are the GHZ family with $\tau^{(3)} = 1$ and all 2-tangles τ vanishing, and the W family with $\tau^{(3)} = 0$ and all 2-tangles equal to 1. Both families are unequivalent under SLOCC transformations [16].

The 3-tangle of the state (6) is $\tau^{(3)} = \frac{1}{4}$ and the reduced density matrices for

$$\text{subsystems } A-B, A-C \text{ and } B-C \text{ are } \rho_{AB} = \frac{1}{4} \begin{pmatrix} 2 & . & 1 & -1 \\ . & . & . & . \\ 1 & . & 1 & -1 \\ -1 & . & -1 & 1 \end{pmatrix}, \rho_{AC} =$$

$$\frac{1}{4} \begin{pmatrix} 1 & -1 & 1 & . \\ -1 & 1 & -1 & . \\ 1 & -1 & 2 & . \\ . & . & . & . \end{pmatrix} \text{ and } \rho_{BC} = \frac{1}{4} \begin{pmatrix} 2 & -1 & -1 & . \\ -1 & 1 & . & . \\ -1 & . & 1 & . \\ . & . & . & . \end{pmatrix}. \text{ The set of eigenvalues of } \rho\tilde{\rho}$$

associated to subsystems $A-B$ and $A-C$ are $\{\frac{1}{16}(3+2\sqrt{2}), \frac{1}{16}(3-2\sqrt{2}), 0, 0\}$ leading to the tangles $\tau_{AB} = \tau_{AC} = \frac{1}{4}$ and the set of eigenvalues of $\rho\tilde{\rho}$ associated to the subsystem $B-C$ is $\{\frac{1}{16}, \frac{1}{16}, 0, 0\}$ leading to a vanishing tangle τ_{BC} .

All states encoded by rows of g_1 and g_2 behaves as state (6). We call a tripartite entanglement characterized by the tangles $\tau^{(3)} = \tau_{AB} = \tau_{AC} = \frac{1}{4}$ and $\tau_{BC} = 0$ an *incomplete balanced entanglement*. Fully balanced tripartite entanglement is found in [1, 11] and in the next section below devoted to the E_8 lattice and other dense lattices.

3. Entanglement arising from Barnes-Wall lattices

The family of Barnes-Wall lattices L_n , of even dimension 2^n , is related to self-dual codes and the real Clifford group. The first four members of the family, for $n = 1$ to 4, are isomorphic to the densest known lattices of the corresponding dimension, i.e. the root lattices A_2 , D_4 , E_8 and the Barnes-Wall lattice BW_{16} .

The automorphism group of L_n (in dimension 2^n , $n \neq 3$) corresponds to the real Clifford group $\mathcal{C}_n^+$ (defined as the normalizer in the general orthogonal group $O(2^n)$ of the generalized Pauli group on n real qubits: the notation comes from [11]). For $n = 3$ qubits, L_3 is E_8 and $\mathcal{C}_3^+$ is the second largest maximal subgroup of $W'(E_8)$, the derived subgroup of the Weyl group $W(E_8)$. In the sequel, we concentrate on the entanglement arising from the lattices embodying two, three and four qubits, respectively.

Two-qubit entanglement and the lattice D_4

Among the several representations of the lattice D_4 , we select the one with basis $M_r = \{(1, 1, 0, 0), (1, -1, 0, 0), (0, 1, -1, 0), (0, 0, 1, -1)\}$, that has the automorphism group

$$\text{Aut}(D_4) = \left\langle \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & 1 \\ 1 & 1 & 1 & -1 \\ 1 & -1 & -1 & -1 \\ -1 & -1 & 1 & -1 \end{pmatrix}, \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & 1 \\ -1 & 1 & 1 & 1 \\ -1 & -1 & 1 & -1 \\ 1 & 1 & 1 & -1 \end{pmatrix} \right\rangle, \quad (8)$$

Going back to (8), it is clear that the rows of the two generators of $\text{Aut}(D_4)$ encode the (entangled) eigenstates of the triple s (as was the case for the generator S). Thus, two-qubit maximal entanglement is at the heart of the symmetries of this representation on the lattice D_4 . The group $\text{Aut}(D_4)$, of order 1152, is (up to isomorphism) the same object than the two-qubit real Clifford group $\mathcal{C}_2^+$ and the Weyl group $W(F_4)$ of the 24-cell.

Three-qubit entanglement and the lattice E_8

Among the several representations of the lattice D_4 , we select the one with basis

$$M = \{(2, -2, 0, 0, 0, 0, 0, 0), (0, 2, -2, 0, 0, 0, 0, 0), (0, 0, 2, -2, 0, 0, 0, 0), (0, 0, 0, 2, -2, 0, 0, 0), \\ (0, 0, 0, 0, 2, -2, 0, 0), (0, 0, 0, 0, 0, 2, -2, 0), (0, 0, 0, 0, 0, 0, 2, -2), (1, 1, 1, 1, 1, -1, -1, -1)\}$$

that has the automorphism group

$$\text{Aut}(E_8) = \langle g_1, g_2 \rangle,$$

$$\begin{aligned}
\text{with } g_1 &= \frac{1}{2} \begin{pmatrix} . & . & -1 & -1 & 1 & . & 1 & . \\ 1 & -1 & . & . & . & -1 & . & -1 \\ . & . & 1 & -1 & 1 & . & -1 & . \\ -1 & -1 & . & . & . & 1 & . & -1 \\ -1 & -1 & . & . & . & -1 & . & 1 \\ . & . & -1 & 1 & 1 & . & -1 & . \\ . & . & -1 & -1 & -1 & . & -1 & . \\ 1 & -1 & . & . & . & 1 & 0 & 1 \end{pmatrix} \\
\text{and } g_2 &= \frac{1}{2} \begin{pmatrix} 1 & -1 & . & . & . & 1 & 1 & . \\ -1 & 1 & . & . & . & 1 & 1 & . \\ . & . & 1 & 1 & 1 & . & . & 1 \\ . & . & -1 & -1 & 1 & . & . & 1 \\ . & . & 1 & -1 & -1 & . & . & 1 \\ -1 & -1 & . & . & . & -1 & 1 & . \\ . & . & -1 & 1 & -1 & . & . & 1 \\ -1 & -1 & . & . & . & 1 & -1 & . \end{pmatrix}.
\end{aligned} \tag{9}$$

The group $\text{Aut}(E_8)$ is of order $4!6!8! = 696\,729\,600$.

Let us consider the state encoded by the first row of g_2

$$|\psi\rangle = \frac{1}{2}(|000\rangle - |001\rangle + |101\rangle + |110\rangle). \tag{10}$$

The tangles attached to the state (10) are calculated, as we did for the state (6) of the previous section. They are such that $\tau^{(3)} = \tau_{AC} = \tau_{AB} = \tau_{BC} = \frac{1}{4}$ corresponding to fully *balanced tripartite entanglement*. The same results holds for all states encoded by the rows of g_1 and g_2 .

One can conclude that this type of entanglement, first featured in [1, 11], is a specific property of the lattice E_8 . As mentioned previously, the 3-qubit real Clifford group $\mathcal{C}_3^+$ of order 2 580 480 is a maximal subgroup of $\text{Aut}'(E_8)$ (the second largest one); the largest maximal subgroup is the automorphism group $\text{Aut}(E_7)$ of the lattice E_7 .

As a second representation of the lattice E_8 , we select the one obtained from the construction “A” on the extended Hamming code [8, 4, 4] [12] §.

$$\begin{aligned}
\text{Aut}(E_8) &= \langle g_1, g_2 \rangle, \\
\text{with } g_1 &= \frac{1}{2} \begin{pmatrix} . & 1 & -1 & . & 1 & . & . & 1 \\ 1 & -1 & . & . & 1 & . & -1 & . \\ -1 & . & -1 & . & . & . & -1 & -1 \\ . & . & . & -1 & . & . & . & . \\ 1 & 1 & . & . & -1 & . & -1 & . \\ . & . & . & . & . & -1 & . & . \\ . & -1 & 1 & . & -1 & . & . & 1 \\ 1 & . & -1 & . & . & . & 1 & -1 \end{pmatrix}
\end{aligned}$$

§ One may use the following commands in Magma [13] “C:=HammingCode(GF(2),3); C:=ExtendCode(C); L:=Lattice(C,”A”); aut:=AutomorphismGroup(L:NaturalAction);aut;”

$$\text{and } g_2 = \frac{1}{2} \begin{pmatrix} \cdot & 1 & \cdot & \cdot & \cdot & 1 & -1 & -1 \\ 1 & \cdot & -1 & -1 & 1 & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & -1 & -1 & \cdot & -1 \\ -1 & -1 & \cdot & -1 & \cdot & \cdot & \cdot & -1 \\ \cdot & \cdot & \cdot & 1 & 1 & \cdot & 1 & -1 \\ \cdot & -1 & -1 & 1 & \cdot & \cdot & -1 & \cdot \\ 1 & -1 & 1 & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & -1 & \cdot & -1 & 1 & 1 & \cdot \end{pmatrix}. \quad (11)$$

Here we get two types of generators. In the generator g_1 , the rows 4 and 6 encode non entangled states while the other rows encode a state such as

$$|\psi\rangle = \frac{1}{2}(|001\rangle - |010\rangle + |100\rangle + |111\rangle), \quad (12)$$

with $\tau^{(3)} = 1$ and vanishing 2-tangles. Such a state is of the GHZ type. The second generator g_2 encodes states of the type (6) corresponding to incomplete balanced entanglement.

Four-qubit entanglement and the lattice Λ_{16}

For four real qubits, the Clifford group is $\mathcal{C}_4^+ \equiv \text{Aut}(\Lambda_{16})$, of order 89 181 388 800, associated to the Barnes-Wall lattice Λ_{16} [7]. We use construction “B” associated to the [16, 5, 8] Reed-Muller code (see [12], p. 141) ||.

One gets two 16×16 gates generating $\text{Aut}(\Lambda_{16})$ with rows encoding entangled states such as the factorizable state

$$\begin{aligned} |\psi\rangle &= \frac{1}{2}(|0000\rangle - |0011\rangle - |0101\rangle + |0110\rangle) \\ &= \frac{1}{2}|0\rangle((|000\rangle - |011\rangle - |101\rangle + |110\rangle)). \end{aligned} \quad (13)$$

It is easy to quantify the residual tripartite entanglement of state (13). Using (7), one gets true tripartite entanglement $\tau^{(3)} = 1$ corresponding to the GHZ family. All states encoded by the rows in the generators of $\text{Aut}(\Lambda_{16})$ display vanishing 4-tangle (the four-tangle is calculated from the expressions given in [15]). Presumably, all residual three-tangles equal unity as for GHZ-type states).

4. Entanglement arising from the D_{12}^+ lattice

The unique indecomposable 12-dimensional unimodular lattice is D_{12}^+ . It can be generated from the construction “A” applied to the [12, 6, 6] extended Golay code over $GF(3)$ ¶.

The automorphism group is of order $2^{21}3^55^27^111$. As for the previous cases, the rows of the generators encode entangled states. If we look at the system as a bipartite

|| in Magma, the code C:=ReedMullerCode(1,4); L:=Lattice(C,“B”);
aut:=AutomorphismGroup(L:NaturalAction);
¶ In Magma, one uses the following commands “C := GolayCode(GF(3), true); L:=Lattice(C,“A”);
aut:=AutomorphismGroup(L:NaturalAction);”

3×4 system, the Schmidt rank⁺ of the generators is found to be three. There are several types of states. We restrict our account to one of them, that we consider as the state of a $3 \times 2 \times 2$ system, i.e. a qutrit coupled to two qubits.

$$|\psi\rangle = (2|000\rangle + |001\rangle - |010\rangle + |011\rangle + |110\rangle + |200\rangle)/3. \quad (14)$$

The reduced density matrix of the two-qubit bipartite system is found to be

$$\rho_{BC} = \frac{1}{9} \begin{pmatrix} 5 & 2 & -2 & 2 \\ 1 & 1 & -1 & 1 \\ -2 & -1 & 2 & -1 \\ 2 & 1 & -1 & 1 \end{pmatrix}$$

The set of eigenvalues of $\rho\tilde{\rho}$ associated the two-qubit subsystem $B - C$ is obtained as $\{\frac{2}{81}(10 + 3\sqrt{11})^{\pm 1/2}, 0, 0\}$, corresponding the two-tangle $\tau_{BC} \sim 0.44^*$.

Then, one can compute the reduced density matrix ρ_{AB} and ρ_{AC} ; they correspond to a qutrit/qubit system. According to the Peres criterion [2]‡, a necessary and sufficient condition for the existence of entanglement between parties A and B is that the partial

⁺ The Schmidt rank can easily be obtained in Magma. Magma provides the Smith normal form for any matrix A with entries in an Euclidean ring (a field K , the ring $\mathbb{Z}$ of integers, the ring $K[x]$ of polynomials with coefficients over a field K ...). Given an $m \times n$ matrix A over an Euclidean ring R , one can find invertible square matrices P and Q such that $PAQ = S$, where S is a diagonal matrix: the Smith normal form of A . This is equivalent to the singular value decomposition $A = P^{-1}SQ^{-1}$. The non-zero diagonal entries of the Smith form of A are unique up to multiplication by a unit of R and are also called elementary divisors. Thus, the Schmidt rank of A equals the rank of S .

^{*} Note that the same row of the selected generator may be thought as attached to a $2 \times 2 \times 3$ system, i.e. two-qubits coupled to a qutrit. In that case, the state has to be rewritten as

$$|\psi\rangle = (2|000\rangle + |001\rangle + |010\rangle - |100\rangle - |200\rangle + |210\rangle)/3. \quad (15)$$

and the new reduced density matrix of the two-qubit bipartite system is found to be

$$\rho_{BC} = \frac{1}{9} \begin{pmatrix} 2 & 2 & 1 & . \\ 2 & 1 & 1 & . \\ 1 & 1 & 2 & . \\ . & . & . & . \end{pmatrix}$$

The set of eigenvalues of $\rho\tilde{\rho}$ associated the two-qubit subsystem $B - C$ is now obtained as $\{\frac{1}{81}(3 \pm 2\sqrt{2}), 0, 0\}$, corresponding the two-tangle $\tau_{BC} \sim 0.050$.

Another selected generator of $\text{Aut}(D_{12}^+)$, seen as a $3 \times 2 \times 2$ system, can easily be found to have a row with entanglement similar to that of (15).

‡ According to the Peres partial transpose (PPT) theorem, a state ρ_{AB} of a bipartite system A-B with dimension $d = d_A d_B \leq 6$ is separable iff

$$(\rho_{AB}^{T_A})_{i\alpha, j\beta} = (\rho_{AB}^{T_A})_{j\alpha, i\beta}$$

has non negative eigenvalues. For $d = d_A d_B > 6$, if the partial transpose matrix $\rho_{AB}^{T_A}$ has negative eigenvalues, then the bipartite system $A - B$ is entangled. But they are states, known as bound entangled states, that satisfy the Peres positivity criterion, but still are entangled.

transpose matrix $\rho_{AB}^{T_A}$ contains at least one negative eigenvalue. A similar criterion holds for the entanglement between parties A and C .

The sets of eigenvalues of matrices

$$\rho_{AB}^{T_A} = \frac{1}{9} \begin{pmatrix} 5 & . & . & 2 & -1 & . \\ . & 2 & 2 & -1 & . & . \\ . & 2 & . & . & . & . \\ . & -1 & . & 1 & . & . \\ 2 & . & . & . & 1 & . \\ -1 & . & . & . & . & . \end{pmatrix}, \quad \rho_{AC}^{T_A} = \frac{1}{9} \begin{pmatrix} 5 & 1 & -1 & 1 & 2 & 1 \\ 1 & 2 & . & . & . & . \\ -1 & . & 1 & . & . & . \\ 1 & . & . & . & . & . \\ 2 & . & . & . & 1 & . \\ 1 & . & . & . & . & . \end{pmatrix}.$$

are $\{0.663, 0.392, 0.092, 0.047, -0.151, -0.0044\}$ and $\{0.717, 0.211, 0.111, 0.047, -0.008\}$, respectively.

These calculations ensure that bipartite entanglement is spread over all parties.

5. Entanglement arising from the Leech lattice

The densest known lattice in dimension 24 is the Leech lattice Λ_{24} with kissing number 196560. It is the unique Euclidean lattice that is even, unimodular and with non-zero vectors at least two. The automorphism group is called $\text{Co}_0 = \mathbb{Z}_2.\text{Co}_1$, where the Conway group Co_1 is sporadic, with order $2^{21}.3^9.5^4.7^2.11.13.23$.

One elegant construction of Λ_{24} makes use of the MOG (Miracle Octad Generator) coordinates, that are obtained from the codewords of the hexacode ([12], p. 132). In this representation, $\text{aut}(\Lambda_{24})$ is entangled. One can see the states encoded by the rows of the two generators as coming from a 6×4 quantum system and for them one finds that the Schmidt rank is four. The amount of entanglement that we find with this approach is quite inhomogeneous. We restrict to a single example, the two-qubit/sextit state

$$\begin{aligned} |\psi\rangle = & (-2|001\rangle + |100\rangle + |101\rangle + |200\rangle + |210\rangle + |300\rangle \\ & + |311\rangle - |500\rangle - |501\rangle - 2|510\rangle)/4, \end{aligned} \quad (16)$$

with reduced density matrix

$$\rho_{BC} = \frac{1}{16} \begin{pmatrix} 4 & 2 & 3 & 1 \\ 2 & 6 & 2 & . \\ 3 & 2 & 5 & . \\ 1 & . & . & 1 \end{pmatrix}$$

and eigenvalues of $\rho\tilde{\rho}$ as $\{\frac{1}{64}(9 \pm 4\sqrt{2}), \frac{1}{256}(3 \pm 2\sqrt{2})\}$. It follows that the two-tangle for this states is $\tau_{BC} \sim 0.00536$.

It can be shown that the partial traces $\rho_{AB}^{T_A}$ and $\rho_{AC}^{T_A}$ violate the Peres positivity criterion so that there also is entanglement between the parties $A - B$ and $A - C$.

It would be desirable to analyze the entanglement in Λ_{24} as arising from a three-qubit/qutrit system but the methods to achieve this goal seem not yet to be available.

6. Conclusion

The symmetry group acting on the generating basis of even Euclidean lattices has been described as a group of entangling quantum gates. Barnes-Wall lattices are known to be related to the Clifford group of quantum error correction [7, 8] and to discretised Hilbert spaces [3] but the occurrence of Leech lattice is novel in this context. It may be that real world systems, like quark systems, or codons, or even black holes [6], exploit dimension $24 = 2^3 \times 3$ as three-qubits and qutrits at the same time. Leech lattice may also be constructed from Leech roots in Lorentzian space $\mathbb{R}^{25,1}$ (see [12], chaps. 26-27) and dimension 26 corresponds to the original version of string theory through the no-ghost theorem. These new vistas (see also [17]), and their connection to tripartite entanglement, will be explored in future work.

Acknowledgements

The author acknowledges Bernd Souvignier for his reading of an earlier version of the manuscript and R. Mosseri for his feedback about the geometry of the discretised Hilbert space. The work started in December 2009 during a visit at de Brún Center for Computational Algebra in Galway under the invitation of Michael Mc Gettrick. It was presented at International Conference on Quantum Optics and Quantum Computing (ICQOQC-11) in Noida and at the 18th Central European Workshop on Quantum Optics (CEWQO 2011) in Madrid.

Bibliography

- [1] Planat M, Levay P and Saniga M 2010 Balanced Tripartite Entanglement, the Alternating Group A_4 and the Lie Algebra $sl(3, C) \oplus u(1)$. *Rep. Math. Phys.* **66** (3) 451-462.
- [2] Horodecki R, Horodecki P, Horodecki M and Horodecki K 2009 Quantum entanglement. *Rev. Mod. Phys.* **81** 865-942.
- [3] Mosseri R and Dandoloff R 2001 Geometry of entangled states, Bloch spheres and Hopf fibrations. *J. Phys. A: Math. Gen.* **34** 10243;
Mosseri R and Ribeiro P 2007 Entanglement and Hilbert space geometry for systems with a few qubits. *Math. Struct. in Comp. Science* **17** 1117.
- [4] Aravind P K, in: R.S. Cohen et al (Eds) 1997 *Potentiality, entanglement and passion-at-a-distance* (Kluwer Academic Publishers) p. 53;
Sugita A 2007 Borromean entanglement revisited. Preprint 0704.1712 (quant-ph).
- [5] Planat M and Saniga M 2008 On the Pauli graphs of N -qudits. *Quant. Inf. & Comp.* **8** 127.
- [6] Levay P, Saniga M and Vrana P 2008 Three-qubit operators, the split Cayley hexagon of order two, and black holes. *Phys. Rev. D* **78** 124022.
- [7] Nebe G, Rains E M and Sloane N J A 2006 *Self-dual codes and invariant theory* (Springer, Berlin).
- [8] Calderbank A R, Rains E M, Schor P W and Sloane N J A 1998 Quantum Error Correction via Codes over $GF(4)$ *IEEE Trans. Inform. Th.* **44** 1369;
Planat M and Solé P 2009 Clifford groups of quantum gates, BN-pairs and smooth cubic surfaces. *J. Phys. A: Math. Theor* **42** 042003.
- [9] Mermin N D 1993 Hidden variables and the two theorems of John Bell. *Rev. Mod. Phys.* **65** 803.
- [10] Planat M, Saniga M and Kibler M R 2006 Quantum entanglement and projective ring geometry. *SIGMA* **2** Paper 066.

- [11] Planat M 2010 Three-qubit entangled embeddings of CPT and Dirac groups within E_8 Weyl group. *Int. J. Theor. Phys.* **49** 1044-1054.
- [12] J. H. Conway and N. J. A. Sloane 1998 *Sphere packings, lattices and groups (third edition)* (Springer New York).
- [13] Bosma W, Cannon J and Playoust C 1997 The Magma algebra system I. The user language *J. Symbolic Comp.* **24** 235;
Plesken W and Souvignier B 1997 Computing isometries of lattices. *J. Symbolic Comp.* **24** 327.
- [14] Harrary F 2000 The automorphism group of a hypercube. *J. Univ. Comp. Sc.* **6** 136.
- [15] Sharma S S and Sharma N K 2010 Four-tangle for pure states. *Phys. Rev. A* **82** 012340.
- [16] Coffman V, Kundu J and Wootters W. K. 2000 Distributed entanglement. *Phys. Rev. A* **61** 052306.
- [17] D. Allcock, On the Y555 complex reflection group, *Journal of Algebra* 322 (2009) 1454.